

ORIGINAL ARTICLE

Whole-genome DNA methylation status associated with clinical PTSD measures of OIF/OEF veterans

R Hammamieh^{1,11}, N Chakraborty^{2,11}, A Gautam^{1,11}, S Muhie^{2,11}, R Yang^{3,11}, D Donohue², R Kumar³, BJ Daigle Jr⁴, Y Zhang⁵, DA Amara⁶, S-A Miller², S Srinivasan², J Flory⁷, R Yehuda⁷, L Petzold⁵, OM Wolkowitz⁸, SH Mellon⁹, L Hood¹⁰, FJ Doyle III⁵, C Marmar⁶ and M Jett¹

Emerging knowledge suggests that post-traumatic stress disorder (PTSD) pathophysiology is linked to the patients' epigenetic changes, but comprehensive studies examining genome-wide methylation have not been performed. In this study, we examined genome-wide DNA methylation in peripheral whole blood in combat veterans with and without PTSD to ascertain differentially methylated probes. Discovery was initially made in a training sample comprising 48 male Operation Enduring Freedom (OEF)/Operation Iraqi Freedom (OIF) veterans with PTSD and 51 age/ethnicity/gender-matched combat-exposed PTSD-negative controls. Agilent whole-genome array detected ~5600 differentially methylated CpG islands (CpGI) annotated to ~2800 differently methylated genes (DMGs). The majority (84.5%) of these CpGIs were hypermethylated in the PTSD cases. Functional analysis was performed using the DMGs encoding the promoter-bound CpGIs to identify networks related to PTSD. The identified networks were further validated by an independent test set comprising 31 PTSD+/29 PTSD- veterans. Targeted bisulfite sequencing was also used to confirm the methylation status of 20 DMGs shown to be highly perturbed in the training set. To improve the statistical power and mitigate the assay bias and batch effects, a union set combining both training and test set was assayed using a different platform from Illumina. The pathways curated from this analysis confirmed 65% of the pool of pathways mined from training and test sets. The results highlight the importance of assay methodology and use of independent samples for discovery and validation of differentially methylated genes mined from whole blood. Nonetheless, the current study demonstrates that several important epigenetically altered networks may distinguish combat-exposed veterans with and without PTSD.

Translational Psychiatry (2017) 7, e1169; doi:10.1038/tp.2017.129; published online 11 July 2017

INTRODUCTION

Adverse life experiences alter the epigenetic profile^{1–3} in a manner that is salient for pathophysiology of post-traumatic stress disorder (PTSD).^{4–6} Changes in methylation status of the glucocorticoid receptor gene have been reported previously in combat veterans with PTSD.⁷ Methylation changes in these same genes were also observed in association with parental trauma, suggesting that such effects may be related to heritable risk profiles.⁸ Consistent claims were presented by *in vivo* studies.^{9,10} Together, these discoveries drive a strong rationale for screening the epigenetic profiles of patients' blood to identify next-generation strategies for PTSD risk factors, diagnostics and experimental therapeutics. A growing body of cohort-based studies has linked the epigenetic changes with PTSD development,^{11–13} mostly focusing on pre-determined targets such as immunity^{14–16} and neuroendocrinology.^{7,8,17,18}

For the present study, strict inclusion–exclusion criteria were used^{19,20} to identify a training set comprising 48 male veterans with PTSD (PTSD+) and 51 age-/ethnicity-/gender-matched controls (PTSD-). Control veterans experienced war trauma but were negative for current and past PTSD (Supplementary

Table S1). An independent test set comprising 31 PTSD+/29 PTSD- veterans was recruited using the same screening protocol.

Enriched by the differentially methylated genes (DMGs), the epigenetically altered networks are linked to nervous systems' development and function, PTSD-associated somatic complications and endocrine signaling. All of these networks mined from the training set were validated by the test set (Table 1). Subsequently, we consolidated the test and training sets to develop a union set and reevaluated the methylation profile using the improved sample size. The result confirmed 65% of the pathways mined from the test and training sets. Going forward, we will consider the methylation profile from this union set as the discovery set to be confirmed in a new validation set, for which subjects are currently being recruited.

MATERIALS AND METHODS

Ethical statement

The Institutional Review Boards of the US Army Medical Research and Materiel Command, the New York University Langone Medical Center (New York, NY, USA), the Icahn School of Medicine at Mt Sinai (New York, NY, USA) and the James J Peters Veterans Administration Medical Center

¹Integrative Systems Biology, US Army Center for Environmental Health Research, Frederick, MD, USA; ²USACEHR, The Geneva Foundation, Frederick, MD, USA; ³Advanced Biomedical Computing Center, Frederick National Laboratory for Cancer Research, Frederick, MD, USA; ⁴Departments of Biological Sciences and Computer Science, The University of Memphis, Memphis, TN, USA; ⁵School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA; ⁶Steven and Alexandra Cohen Veterans Center for the Study of Posttraumatic Stress and Traumatic Brain Injury, Department of Psychiatry, NYU School of Medicine, New York, NY, USA; ⁷Department of Psychiatry, Mount Sinai School of Medicine, James J Peters Veterans Administration Medical Center, Bronx, NY, USA; ⁸Department of Psychiatry, University of California San Francisco, San Francisco, CA, USA; ⁹Department of Ob/Gyn, Reproductive Sciences, University of California, San Francisco, San Francisco, CA, USA and ¹⁰Institute for Systems Biology, Seattle, WA, USA. Correspondence: Dr R Hammamieh, Integrative Systems Biology, US Army Center for Environmental Health Research, 568 Doughten Drive, Fort Detrick, Frederick, MD 21702-5010, USA. E-mail: Rasha.Hammamieh1.civ@mail.mil

¹¹These authors contributed equally to this work.

Received 3 May 2017; accepted 4 May 2017

REPORT DOCUMENTATION PAGE					<i>Form Approved</i> OMB No. 0704-0188							
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.												
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.												
1. REPORT DATE (DD-MM-YYYY) 11-07-2017		2. REPORT TYPE Research			3. DATES COVERED (From - To) 01-2011 to 05-2017							
4. TITLE AND SUBTITLE Whole-genome DNA methylation status associated with clinical PTSD measures of OIF/OEF veterans				5a. CONTRACT NUMBER								
				5b. GRANT NUMBER								
				5c. PROGRAM ELEMENT NUMBER ISB								
				5d. PROJECT NUMBER								
				5e. TASK NUMBER								
6. AUTHOR(S) Hammamieh, Rasha; Chakraborty, Nabarun; Gautam, Aarti; Muhie, Seid; Yang, Routing; Donohue, Duncan; Kumar, Raina; Daigle, BJ, Jr.; Zhang, Y; Amara, DA; Miller, Stacy-Ann; Srinivasan, Seshamalini; Flory, Janine; Yehuda, Rachel; Petzold, L; Wolkowitz, OH; Mellon, SH; Hood, L; Doyle FJ, III; Marmar, C; Jett, Marti				5f. WORK UNIT NUMBER								
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S Army Center for Environmental Health Research 568 Doughten Drive Fort Detrick, MD 21702-5010					8. PERFORMING ORGANIZATION REPORT NUMBER 16-017							
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical and Materiel Command 504 Scott Street Fort Detrick, MD 21702-5010					10. SPONSOR/MONITOR'S ACRONYM(S) USAMRMC							
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)							
12. DISTRIBUTION/AVAILABILITY STATEMENT public release												
13. SUPPLEMENTARY NOTES Published in Translational Psychiatry (2017) 7, e1169												
14. ABSTRACT Emerging knowledge suggests that post-traumatic stress disorder (PTSD) pathophysiology is linked to the patients' epigenetic changes, but comprehensive studies examining genome-wide methylation have not been performed. In this study, we examined genome-wide DNA methylation in peripheral whole blood in combat veterans with and without PTSD to ascertain differentially methylated probes. Discovery was initially made in a training sample comprising 48 male Operation Enduring Freedom (OEF)/Operation Iraqi Freedom (OIF) veterans with PTSD and 51 age/ethnicity/gender-matched combat-exposed PTSD-negative controls. Agilent whole-genome array detected ~ 5600 differentially methylated CpG islands (CpGI) annotated to ~ 2800 differently methylated genes (DMGs). The majority (84.5%) of these CpGIs were hypermethylated in the PTSD cases. Functional analysis was performed using the DMGs encoding the promoter-bound CpGIs to identify networks related to PTSD. The identified networks were further validated by an independent test set comprising 31 PTSD+/29 PTSD - veterans. Targeted bisulfite sequencing was												
15. SUBJECT TERMS												
16. SECURITY CLASSIFICATION OF: <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%; padding: 2px;">a. REPORT</td> <td style="width: 33%; padding: 2px;">b. ABSTRACT</td> <td style="width: 33%; padding: 2px;">c. THIS PAGE</td> </tr> <tr> <td style="text-align: center; padding: 2px;">U</td> <td style="text-align: center; padding: 2px;">U</td> <td style="text-align: center; padding: 2px;">U</td> </tr> </table>			a. REPORT	b. ABSTRACT	c. THIS PAGE	U	U	U	17. LIMITATION OF ABSTRACT SAR		18. NUMBER OF PAGES 7	
a. REPORT	b. ABSTRACT	c. THIS PAGE										
U	U	U										
			19a. NAME OF RESPONSIBLE PERSON Linda M. Brennan									
			19b. TELEPHONE NUMBER (Include area code) 301-619-7530									

Reset

Table 1. The pathways of interest and their status of validation

Pathway	Number of genes	
	Training set	Test set
<i>Nervous system functions</i>		
Addiction	25	15
Aggressive behavior	14	8
Fear response	48	16
Amygdala fear response		
Fear memory consolidation		
Fear memory extinction		
Fear memory potentiation		
Fear-potentiated anxiety		
Fear-potentiated startle		
Long-term impact on the brain	78	18
Long-term fear memory		
Long-term memory		
Long-term synaptic depression		
Long-term synaptic potentiation		
Depression	10	11
Learning	27	9
Associative learning		
Learning or memory		
Traumatic fear learning		
Social withdrawal	10	3
<i>PTSD-associated somatic complications</i>		
Diabetes and insulin signaling	47	19
Premature aging	50	38
Metabolic syndrome		
Telomere management		
Mitochondrial dysfunction		
Inflammation	81	70
Circadian rhythm	40	12
REM sleep	12	2
<i>PTSD-relevant endocrine signaling networks</i>		
Corticotrophin-releasing hormone network	30	16
Dopamine and serotonin signaling	27	12
Glucocorticoid signaling	52	36
HPA axis	28	10
<i>Nervous system development</i>		
Axon guidance	106	42
Cannabinoid management	28	4
Hippocampus development	7	45
Neurogenesis	60	215
Nerve impulse	42	31
Synaptic plasticity	48	74
Abbreviations: HPA, hypothalamus–pituitary–adrenal; REM: rapid eye movement; PTSD, post-traumatic stress disorder.		

(Bronx, NY, USA) approved this study. Study participants gave written and informed consent to participate. The study was conducted in accordance with the provisions of the Helsinki Declaration.

Cohort recruitment and analysis

The recruitment process involved several steps detailed in the Supplementary Table S1 and in previous communications.^{19,20} The training set of 48 PTSD+/51 PTSD– and the test set of 31 PTSD+/29 PTSD– veterans was probed by whole-genome arrays (Agilent, Santa Clara, CA, USA) containing ~27k CpGs. The outcome was normalized to minimize the confounding factors attributed to batch processing.²¹ Functional analysis was performed using those DMGs, which encoded CpGs meeting the cutoff false discovery rate < 0.1.

Next, we merged the training and test sets to develop a union set comprising 79 PTSD+/80 PTSD– veterans, which was probed by whole-genome arrays (Illumina, San Diego, CA, USA) containing 450 k probes. The outcome was corrected to minimize heterogeneous cell populations²² and

age effects, and was screened at $P < 0.05$ to find DMGs. Available GEO databases are as follows: GSE76401 and GSE85399. ClueGo v2.1.2 and Ingenuity pathway analysis were used for network construction, and pathways that we report met the cutoff of $P < 0.05$.

RESULTS

The primary purpose of the present communication was to identify the functional networks associated with combat-related PTSD, and thereby to provide a better understanding of PTSD pathophysiology. To meet this goal, we recruited 48 PTSD+/51 PTSD– veterans as a training set and 31 PTSD+/29 PTSD– veterans as a test set. To increase the statistical power and to minimize any bias of the Agilent high-throughput array platform, we took two measures. First, we constructed a union set by consolidating the training and test sets, following a recently published strategy.^{19,20} Second, we retested the methylation profile, probing the union set using a different array platform manufactured by Illumina. Furthermore, this union set retains sufficient statistical power. Taking a moderate estimate of 50% s. d.'s in probe signals and a relatively conservative estimate for the mean difference (that is, top 1%), 76 people per group should give 95% power to detect an individual probe with a (Bonferroni-adjusted) genome-wide significance of $P < 1.162931e - 07$.

Functional analysis of the training set found a host of PTSD-related networks

In the investigation of the 48 PTSD+/51 PTSD– training set, we identified 5578 differentially methylated CpGs annotated to 3662 genes. We collectively defined the 1698 promoter-bound CpGs and 157 additional divergent promoter regions as the *promoter* regions (Supplementary Figure S4A). Altogether, 4721 CpGs annotated 2401 DMGs that displayed a \log_2 ratio > 0.1 and were defined as hypermethylated. Conversely, 857 CpGs (672 DMGs) displaying a \log_2 ratio < 0.1 were defined as hypomethylated. The remaining DMGs co-enriched by both hyper- and hypomethylated CpGs were excluded from the subsequent functional analysis. For the functional analysis, we used those DMGs, which encoded promoter-bound CpGs, estimated as nearly 60% of total DMGs. Significantly enriched networks with similar functional purposes were grouped together, resulting in four network clusters (Figure 1): nervous system functions (Figure 2a), PTSD-associated somatic complications (Figure 2b), PTSD-relevant endocrine signaling networks (Supplementary Figure S6A) and nervous system development (Supplementary Figure S6B).

Test set validated all the networks identified by the training set. There was a significant ($P < 0.001$) overlap at the DMG level between the 48 PTSD+/51 PTSD– training set and the 31 PTSD+/29 PTSD– test set with 779 DMGs in common between the two sets assayed by the Agilent whole-genome array. Furthermore, a significant agreement was noted at the functional level as all of the networks mined from the training set emerged significantly enriched by DMGs identified from the test set (Table 1).

Union set probed by a different array platform validated a majority of networks identified by the training and test sets

The union set probed by the Illumina array resulted in 3339 DMG, 74.4% of which encoded hypermethylated CpGs (Supplementary Figures S4B and C). One hundred ninety-one DMGs were in common between the training set and union set, and 107 DMGs were in common between the test set and union set (Supplementary Figure S5). There were 852 DMGs encoding promoter-bound CpGs enriched in networks linked to addiction, long-term impact on cerebral functions, social withdrawal, diabetes, aging, inflammation, circadian rhythm, dopamine-




Figure 1. Functional enrichment analysis. In all, 352 DMGs encoding promoter-bound differentially methylated CpGs curated from the training set were enriched for four functional clusters: PTSD-associated somatic complications, PTSD-relevant endocrine signaling, nervous system development and nervous system functions. These clusters were designed to group networks with overlapping functionality. All of these networks were validated by the test set. CRH, corticotrophin-releasing hormone; DMG, differentially methylated gene; GC, glucocorticoid; HPA, hypothalamus–pituitary–adrenal; PTSD, post-traumatic stress disorder; REM, rapid eye movement.

serotonin signaling, neurogenesis, cannabinoid signaling, nerve impulse and synaptic plasticity. In addition, 407 DMGs in shelf and shore regions were enriched in networks associated with REM sleep, circadian rhythm, inflammation, hypothalamic–pituitary–adrenal axis and axon guidance. Altogether, the union set confirmed 15 out of 23 networks mined from the training set and validated by test set. All of the networks clustered under PTSD-associated somatic complications and nervous systems' development were confirmed by the training, test and union sets.

Methylation status of selected DMGs validated by targeted bisulfite sequencing

Forty-two DMGs were selected from the training set based on their methylation status and their relevance to PTSD. Their methylation status was verified by targeted bisulfite sequencing (Zymo Research, Irvine, CA, USA; Table 2).^{23,24} Twenty genes out of forty-two DMGs were confirmed with the Agilent array data. Table 2 lists these genes and their relevance to PTSD and associated comorbidities.

DISCUSSION

Clinical measures were in agreement with the epigenetically altered networks and DMGs

Self-reported clinical measures indicated that veterans with PTSD were concurrently experiencing higher levels of fear, social withdrawal, anxiety, hostility, depression and anger than were controls. Epigenetic investigation of DNA extracted from whole blood revealed networks relevant to these PTSD-associated negative emotions. Greater waist size, waist-to-hip ratio and body mass index¹⁹ were found in PTSD cases as compared with controls and are consistent with the observed pathways associated with

cardiac diseases, diabetes and metabolic syndrome. PTSD-associated immune dysregulation has been previously reported in epigenetic studies.^{14,15,19} Consistent with previous findings,¹⁴ our results found a host of innate immunity-associated genes, consisting of 60% of the entire set of DMGs found altered in PTSD patients. In extending this knowledge, we functionally linked a majority of these genes to mobilization of phagocytic macrophages and leukocytes.

In addition, we identified epigenetically altered networks linked to learning and memory that are relevant for PTSD-associated neurocognitive impairment. Previous epidemiology studies suggested that there was an increased risk of premature aging in PTSD.^{34–36} We identified two epigenetically altered networks relevant to aging. The first network is telomere management and interaction with pathways of two mediators, wnt/ β -catenin³⁷ and p53.³⁸ The epigenetic profile of these aging markers³⁵ was altered in PTSD. The second network is mitochondrial dysfunction, also epigenetically altered in PTSD veterans. Consistent with these markers of premature aging, we found evidence recently for decreased mitochondrial DNA copy numbers in PTSD veterans from this cohort, suggesting a role for energy deprivation in PTSD that escalates the aging process.³⁹

Premature aging^{40,41} and other PTSD-associated somatic complications, such as dysregulation of immunity,⁴² are known to be associated with circadian rhythm. Veterans with PTSD showed epigenetic regulation of some of the key molecular nodes responsible for setting the circadian clock. We identified DMGs encoding CREB3 and GRIN2A, which control photoreception,⁴³ and that are involved in signaling to entrain the circadian clock regulation by *CLOCK* and *PER1* genes.⁴⁴

Epigenetic changes in neurogenic functional pathways were captured by the differential methylation of members of the neural helix–loop–helix family, including *NEUROG1* and *HES1* and their




Figure 2. (a) Network cluster annotated to nervous system functions significantly enriched by DMGs in the training set. (b) Network cluster annotated to PTSD-associated somatic complications significantly enriched by DMGs in the training set. (c) Network cluster annotated to PTSD-relevant endocrine networks significantly enriched by DMGs in the training set. (d) Network cluster annotated to nervous system development networks significantly enriched by DMGs in the training set. In all the figure, red and green circles are hypermethylated and hypomethylated genes, respectively. Sizes of the circles labeled by the annotation terms are correlated with their significance of enrichment. DMG, differentially methylated gene; PTSD, post-traumatic stress disorder.

regulators ATOH-1, Pax6 and NKX2-2.^{45,46} Epigenetic perturbations of networks related to the hypothalamic–pituitary–adrenal axis functions and the synthesis of key feedback regulators, such as corticotrophin and glucocorticoid, as well as epigenetic changes in the serotonergic and dopaminergic networks, may serve as targets for novel therapeutics for PTSD.⁴⁷

Strengths, limitations and future work

The Diagnostic and Statistical Manual of Mental Disorders-IV diagnostic criteria⁴⁸ were used to determine the PTSD status, an approach to clinical phenotyping, which has limitations.

We attempted to maximize signal detection by employing stringent selection criteria including a requirement of Clinician-Administered PTSD scale scores of 40 or greater for PTSD cases and scores less than 20 for controls.^{19,20} Our array-based approach selected two platforms that ensured extensive coverage of the genome and instilled higher confidence in the outcome. We also focused primarily on the promoter regions, as the methylation shifts near transcription start site are most likely to be associated with long-term gene silencing.⁴⁹

Given the biological heterogeneity of PTSD, our findings are limited by the sizes of our discovery, test and union sets.⁵⁰

Table 2. Differentially methylated genes validated by targeted sequencing

Gene symbol	CpG location	Methylation Status	Brief recent literature review (human studies)
AKT1	Chr14: 105262368 (TSS – 287) 105262438 (TSS – 357) 105262494 (TSS – 413)	↑	Associated network is vulnerable to stress-induced anxiety and depression, major comorbidities of PTSD ^{25,26}
BDNF	Chr11: 27744245 (TSS – 639)	↑	BDNF expression was high in human PTSD serum ²⁷ and low in PTSD plasma samples. ²⁸ However, the plasma result was not validated in a subsequent study ²⁹
CNR1	Chr6: 88876636 (TSS – 868) 88876636 (TSS – 1067)	↑	PTSD is significantly associated with SNP haplotype (for C-A and C-G) of CNR1 ³⁰
CREB1	Chr2: 208394337 (TSS+277)	↑	Altered the gene expressions of CREB family occurred in PTSD patients' monocytes ³¹
DMRTA2	Chr1: 50890130 (TSS – 1010)	↓	—
EFS	Chr14: 23835035 (TSS – 192)	↑	—
ELK1	ChrX: 47510240 (TSS – 236)	↓	—
ETS-2	Chr21: 40177278 (TSS+47) 40177531 (TSS+222)	↑	ETS-2 gene family is responsible for growth control, transformation and developmental programs that influence telomere shift and premature aging. Both complications are PTSD-associated ³²
GATA3	Chr10: 8096093 (TSS+226876)	↓	—
HES4	Chr1: 936030 (TSS – 477) 936301 (TSS – 748)	↑	—
LHX1	Chr17: 35292083 (TSS+2687)	↑	—
MET	Chr7: 116311962 (TSS+495) 116312201 (TSS+256)	↑	—
NFATC4	Chr14 24836169 (TSS+24) 24836183 (TSS+38) 24836217 (TSS+72)	↓	This is an immune-associated gene
NR2E1	Chr6: 198486189 (TSS+1024) 198486237 (TSS+976)	↑	NR2E1 deletion produces a highly aggressive phenotype ³³
PAX5	Chr9: 37036906 (TSS – 2429)	↑	—
PDGFB	Chr22: 39638278 (TSS – 1363) 39638353 (TSS – 1438) 39638364 (TSS – 1449)	↑	—
PSD	Chr10: 104178908 (TSS – 6) 104178910 (TSS – 8) 104178916 (TSS – 14) 104178930 (TSS – 28) 104178932 (TSS – 30)	↑	—
PTTG1IP	Chr21: 46294077 (TSS – 258)	↓	—
TRERF1	Chr6: 42420444 (TSS – 660)	↑	—

Abbreviations: ↑, hypermethylated; ↓, hypomethylated; PTSD, post-traumatic stress disorder; SNP, single-nucleotide polymorphism; TSS, transcription start site. (i) Whole-genome array from Agilent, probing of the training set (48/51 PTSD+/–); (ii) targeted bisulfite-sequencing assay (Zymo Research) probing of the training set (48/51 PTSD+/–).

The selection of the Illumina platform was driven by the following three factors: (i) this platform offered nearly twice the number of CpGs to test in comparison to the Agilent platform; (ii) the significantly lower amount of input DNA required for the Illumina assay (500 ng DNA versus 5 µg for the Agilent, assay) satisfied our need to conserve gradually decreasing DNA stocks; and (iii) the growing preference for the Illumina assay in the epigenetics literature^{11,51} was convincing for its selection. The present study recruited the largest cohort size used to date to study the PTSD pathophysiology. The statistical analysis has moderate statistical power attributed to the sample size, which was further enhanced by the strict regulations applied by the pathway enrichment analysis. The epigenetic contributions of many of those genes discovered have been reported as linked to PTSD via transcriptional variations. In addition, many novel epigenetic markers

linked to PTSD were presented here. Together, this study revealed some of the key aspects of PTSD, such as its long-term health implications, which could be best explained by the epigenetic model. However, it is challenging to draw robust mechanistic conclusions due to the non-longitudinal nature of the study; hence, there is a limited scope for making inferences about whether these epigenetic alterations are causes of or consequences of PTSD. This study is also lacking in prospective design, gender balance and systems-wide integration. The findings are compromised further by the fact that the array platforms are potentially unable to provide the extensive coverage typical of deep sequencing.

On the basis of these findings, future work should focus on those epigenetically altered networks presented herein, which showed clinical relevance to PTSD pathophysiology. Our study

presented a knowledge-driven data-mining architecture particularly useful to identify potential biomarkers for a multifactorial disease such as PTSD. In particular, we demonstrated how to use the clinical and physical dimensions as the successful guiding cue to mine the molecular markers linked to disease pathophysiology. This data-mining approach will be practised further in our future study that will recruit a new validation set to confirm the results obtained from the union set serving as the better-powered discovery set. We will also recruit a cohort of female veterans to minimize gender bias. Additional data from blood counts and magnetic resonance imaging will be included. System-wide knowledge integration will be performed to identify PTSD biomarkers with the highest efficacy.^{52–57}

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

The project is supported by the USAMRMC Military Operational Medicine Research Program (MOMRP)/Defense Health Agency (DHA)/Congressional Special Interests (CSI) and MOMRP 190040.

DISCLAIMER

The views, opinions and/or findings contained in this report are those of the authors and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other official documentation.

REFERENCES

- Uher R, Weaver IC. Epigenetic traces of childhood maltreatment in peripheral blood: a new strategy to explore gene-environment interactions. *Br J Psychiatry* 2014; **204**: 3–5.
- Tammen SA, Friso S, Choi SW. Epigenetics: the link between nature and nurture. *Mol Aspects Med* 2013; **34**: 753–764.
- Feil R, Fraga MF. Epigenetics and the environment: emerging patterns and implications. *Nat Rev Genet* 2011; **13**: 97–109.
- Chakraborty N, Meyerhoff J, Gautama A, Muhie A, Jibitua M, De Limab TCM et al. Gene and stress history interplay in emergence of PTSD-like features. *Behav Brain Res* 2015; **292**: 266–277.
- Spiegel D. An ingenious study of intergenerational transmission of the effects of PTSD. *Am J Psychiatry* 2014; **171**: 811–813.
- Stein MB, Jang KL, Taylor S, Vernon PA, Livesley WJ. Genetic and environmental influences on trauma exposure and posttraumatic stress disorder symptoms: a twin study. *Am J Psychiatry* 2002; **159**: 1675–1681.
- Yehuda R, Flory JD, Bierer LM, Henn-Haase C, Lehrner A, Desarnaud F et al. Lower methylation of glucocorticoid receptor gene promoter 1F in peripheral blood of veterans with posttraumatic stress disorder. *Biol Psychiatry* 2015; **77**: 356–364.
- Yehuda R, Daskalakis NP, Lehrner A, Desarnaud F, Bader HN, Makotkine I et al. Influences of maternal and paternal PTSD on epigenetic regulation of the glucocorticoid receptor gene in Holocaust survivor offspring. *Am J Psychiatry* 2014; **171**: 872–880.
- Zovick IB, Sweatt JD. Epigenetic mechanisms in learned fear: implications for PTSD. *Neuropsychopharmacology* 2013; **38**: 77–93.
- Pizzimenti CL, Lattal KM. Epigenetics and memory: causes, consequences and treatments for post-traumatic stress disorder and addiction. *Genes Brain Behav* 2015; **14**: 73–84.
- Norholm SD, Jovanovic T, Smith AK, Binder E, Klengel T, Conneely K et al. Differential genetic and epigenetic regulation of catechol-O-methyltransferase is associated with impaired fear inhibition in posttraumatic stress disorder. *Front Behav Neurosci* 2013; **7**: 30.
- Uddin M, Galea S, Chang SC, Koenen KC, Goldmann E, Wildman DE et al. Epigenetic signatures may explain the relationship between socioeconomic position and risk of mental illness: preliminary findings from an urban community-based sample. *Biodemogr Soc Biol* 2013; **59**: 68–84.
- Rusiecki JA, Chen L, Srikantan V, Zhang L, Yan L, Polin ML et al. DNA methylation in repetitive elements and post-traumatic stress disorder: a case-control study of US military service members. *Epigenomics* 2012; **4**: 29–40.

- Breen MS, Maihofer AX, Glatt SJ, Tylee DS, Chandler SD, Tsuang MT et al. Gene networks specific for innate immunity define post-traumatic stress disorder. *Mol Psychiatry* 2015; **20**: 1538–1545.
- Uddin M, Aiello AE, Wildman DE, Koenen KC, Pawelec G, de Los Santos R et al. Epigenetic and immune function profiles associated with posttraumatic stress disorder. *Proc Natl Acad Sci USA* 2010; **107**: 9470–9475.
- Rusiecki JA, Byrne C, Galdzicki Z, Srikantan V, Chen L, Poulin M et al. PTSD and DNA methylation in select immune function gene promoter regions: a repeated measures case-control study of U.S. military service members. *Front Psychiatry* 2013; **4**: 56.
- Vukojevic V, Kolassa IT, Fastenrath M, Gschwind L, Spalek K, Milnik A et al. Epigenetic modification of the glucocorticoid receptor gene is linked to traumatic memory and post-traumatic stress disorder risk in genocide survivors. *J Neurosci* 2014; **34**: 10274–10284.
- Labonte B, Azoulay N, Yerko V, Turecki G, Brunet A. Epigenetic modulation of glucocorticoid receptors in posttraumatic stress disorder. *Transl Psychiatry* 2014; **4**: e368.
- Lindqvist D, Wolkowitz OM, Mellon S, Yehuda R, Flory JD, Henn-Haase C et al. Proinflammatory milieu in combat-related PTSD is independent of depression and early life stress. *Brain Behav Immun* 2014; **42**: 81–88.
- Yan X, Brown AD, Lazar M, Cressman VL, Henn-Haase C, Neylan TC et al. Spontaneous brain activity in combat related PTSD. *Neurosci Lett* 2013; **547**: 1–5.
- Leek JT, Scharpf RB, Bravo HC, Simcha D, Langmead B, Johnson WE et al. Tackling the widespread and critical impact of batch effects in high-throughput data. *Nat Rev Genet* 2010; **11**: 733–739.
- Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH et al. DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC Bioinformatics* 2012; **13**: 86.
- Wilmot B, Fry R, Smeester L, Musser ED, Mill J, Nigg JT. Methylomic analysis of salivary DNA in childhood ADHD identifies altered DNA methylation in VIPR2. *J Child Psychol Psychiatr* 2016; **57**: 152–160.
- Ashktorab H, Daremipouran M, Goel A, Varma S, Leavitt R, Sun X et al. DNA methylome profiling identifies novel methylated genes in African American patients with colorectal neoplasia. *Epigenetics* 2014; **9**: 503–512.
- Krishnan V, Han MH, Mazei-Robison M, Iniguez SD, Ables JL, Vialou V et al. AKT signaling within the ventral tegmental area regulates cellular and behavioral responses to stressful stimuli. *Biol Psychiatry* 2008; **64**: 691–700.
- Dahlhoff M, Siegmund A, Golub Y, Wolf E, Holsboer F, Wotjak CT. AKT/GSK-3 β /beta-catenin signalling within hippocampus and amygdala reflects genetically determined differences in posttraumatic stress disorder like symptoms. *Neuroscience* 2010; **169**: 1216–1226.
- Martinotti G, Sepede G, Brunetti M, Ricci V, Gambi F, Chillemi E et al. BDNF concentration and impulsiveness level in post-traumatic stress disorder. *Psychiatr Res* 2015; **229**: 814–818.
- Dell'Osso L, Carmassi C, Del Debbio A, Catena Dell'Osso M, Bianchi C, da Pozzo E et al. Brain-derived neurotrophic factor plasma levels in patients suffering from post-traumatic stress disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 2009; **33**: 899–902.
- Su S, Xiao Z, Lin Z, Qiu Y, Jin Y, Wang Z. Plasma brain-derived neurotrophic factor levels in patients suffering from post-traumatic stress disorder. *Psychiatr Res* 2015; **229**: 365–369.
- Lu AT, Ogdie MN, Jarvelin MR, Moilanen IK, Loo SK, McCracken JT et al. Association of the cannabinoid receptor gene (CNR1) with ADHD and post-traumatic stress disorder. *Am J Med Genet B Neuropsychiatr Genet* 2008; **147B**: 1488–1494.
- O'Donovan A, Sun B, Cole S, Rempel H, Lenoci M, Pulliam L et al. Transcriptional control of monocyte gene expression in post-traumatic stress disorder. *Dis Markers* 2011; **30**: 123–132.
- Rameil P, Lecine P, Ghysdael J, Gouilleux F, Kahn-Perles B, Imbert J. IL-2 and long-term T cell activation induce physical and functional interaction between STAT5 and ETS transcription factors in human T cells. *Oncogene* 2000; **19**: 2086–2097.
- Young KA, Berry ML, Mahaffey CL, Saionz JR, Hawes NL, Chang B et al. Fierce: a new mouse deletion of Nr2e1; violent behaviour and ocular abnormalities are background-dependent. *Behav Brain Res* 2002; **132**: 145–158.
- Zhang L, Hu XZ, Li X, Li H, Smerin S, Russell D et al. Telomere length - a cellular aging marker for depression and post-traumatic stress disorder. *Med Hypotheses* 2014; **83**: 182–185.
- Boks MP, van Mierlo HC, Rutten BP, Radstake TR, De Witte L, Geuze E et al. Longitudinal changes of telomere length and epigenetic age related to traumatic stress and post-traumatic stress disorder. *Psychoneuroendocrinology* 2015; **51**: 506–512.
- Lohr JB, Palmer BW, Eidt CA, Ailaboyina S, Mausbach BT, Wolkowitz OM et al. Is post-traumatic stress disorder associated with premature senescence? A review of the literature. *Am J Geriatr Psychiatry* 2015; **23**: 709–725.
- Zhang DY, Wang HJ, Tan YZ. Wnt/beta-catenin signaling induces the aging of mesenchymal stem cells through the DNA damage response and the p53/p21 pathway. *PLoS ONE* 2011; **6**: e21397.

- 38 Artandi SE, Attardi LD. Pathways connecting telomeres and p53 in senescence, apoptosis, and cancer. *Biochem Biophys Res Commun* 2005; **331**: 881–890.
- 39 Bersani FS, Morley C, Lindqvist D, Epel ES, Picard M, Yehuda R. Mitochondrial DNA copy number is reduced in male combat veterans with PTSD. *Progr Neuro-Psychopharmacol Biol Psychiatry* 2015; **64**: 17.
- 40 Kondratov RV, Kondratova AA, Gorbacheva VY, Vykhovanets OV, Antoch MP. Early aging and age-related pathologies in mice deficient in BMAL1, the core component of the circadian clock. *Genes Dev* 2006; **20**: 1868–1873.
- 41 Monk TH. Aging human circadian rhythms: conventional wisdom may not always be right. *J Biol Rhythms* 2005; **20**: 366–374.
- 42 Scheiermann C, Kunisaki Y, Frenette PS. Circadian control of the immune system. *Nat Rev Immunol* 2013; **13**: 190–198.
- 43 Travnickova-Bendova Z, Cermakian N, Reppert SM, Sassone-Corsi P. Bimodal regulation of mPeriod promoters by CREB-dependent signaling and CLOCK/BMAL1 activity. *Proc Natl Acad Sci USA* 2002; **99**: 7728–7733.
- 44 Ko CH, Takahashi JS. Molecular components of the mammalian circadian clock. *Hum Mol Genet* 2006; **15**(Spec No 2): R271–R277.
- 45 Sugimori M, Nagao M, Bertrand N, Parras CM, Guillemot F, Nakafuku M. Combinatorial actions of patterning and HLH transcription factors in the spatiotemporal control of neurogenesis and gliogenesis in the developing spinal cord. *Development* 2007; **134**: 1617–1629.
- 46 Mulvaney J, Dabdoub A. Atoh1, an essential transcription factor in neurogenesis and intestinal and inner ear development: function, regulation, and context dependency. *J Assoc Res Otolaryngol* 2012; **13**: 281–293.
- 47 Asnis GM, Kohn SR, Henderson M, Brown NL. SSRIs versus non-SSRIs in post-traumatic stress disorder: an update with recommendations. *Drugs* 2004; **64**: 383–404.
- 48 Association AP. *Diagnostic Criteria from DSM-IV*. The Association: Washington, DC, 1994.
- 49 Jones PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat Rev Genet* 2012; **13**: 484–492.
- 50 Shao L, Fan X, Cheng N, Wu L, Cheng Y. Determination of minimum training sample size for microarray-based cancer outcome prediction—an empirical assessment. *PLoS ONE* 2013; **8**: e68579.
- 51 Mehta D, Klengel T, Conneely KN, Smith AK, Altmann A, Pace TW et al. Childhood maltreatment is associated with distinct genomic and epigenetic profiles in posttraumatic stress disorder. *Proc Natl Acad Sci USA* 2013; **110**: 8302–8307.
- 52 Gates MA, Holowka DW, Vasterling JJ, Keane TM, Marx BP, Rosen RC. Posttraumatic stress disorder in veterans and military personnel: epidemiology, screening, and case recognition. *Psychol Services* 2012; **9**: 361–382.
- 53 Hayes JP, Vanelzakker MB, Shin LM. Emotion and cognition interactions in PTSD: a review of neurocognitive and neuroimaging studies. *Front Integr Neurosci* 2012; **6**: 89.
- 54 Solomon Z, Helvitz H, Zerach G. Subjective age, PTSD and physical health among war veterans. *Aging Ment Health* 2009; **13**: 405–413.
- 55 Mehta D, Binder EB. Gene x environment vulnerability factors for PTSD: the HPA-axis. *Neuropharmacology* 2012; **62**: 654–662.
- 56 McDonald SD, Calhoun PS. The diagnostic accuracy of the PTSD checklist: a critical review. *Clin Psychol Rev* 2010; **30**: 976–987.
- 57 Thakur Gunjan S, Daigle Jr Bernie J, Dean Kelsey R, Zhang Yuanyang, Rodriguez-Fernandez Maria, Hammamieh Rasha, Yang Ruoting, Jett Marti, Palma Joseph, Petzold Linda R, Doyle III Francis J. Systems biology approach to understanding post-traumatic stress disorder. *Mol. Biosyst* 2015; **11** (4): 980–993. 10.1039/C4MB00404C.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>

© The Author(s) 2017

Supplementary Information accompanies the paper on the Translational Psychiatry website (<http://www.nature.com/tp>)